

Supplementary Online Content

Simon GE, Stewart C, Yarborough BJ, et al. Mortality rates after the first diagnosis of psychotic disorder in adolescents and young adults. *JAMA Psychiatry*. Published online January 31, 2018. doi:10.1001/jamapsychiatry.2017.4437

eTable. Sensitivity Analyses Comparing Mortality Rates With Case and Control Samples Limited to Those With at Least 24 and 36 Months of Prior Enrollment

This supplementary material has been provided by the authors to give readers additional information about their work.

eTable. Sensitivity Analyses Comparing Mortality Rates With Case and Control Samples Limited to Those With at Least 24 and 36 Months of Prior Enrollment

	1 st Diagnosis of Psychotic Disorder		1st Diagnosis of Unipolar Depression		General outpatient	
	No.	Rate per 10,000 per year (95% CI)	No.	Rate per 10,000 per year (95% CI)	No.	Rate per 10,000 per year (95% CI)
ENROLLED >= 12 MONTHS		(n=11,713)		(n=23,415)		(n=35,576)
Total deaths	64	54.6 (41.3, 68.0)	48	20.5 (14.7, 26.3)	24	6.7 (4.0, 9.4)
Self-inflicted Injury/Poisoning	22	18.8 (10.9, 26.6)	26	11.1 (6.8, 15.4)	2	0.6 (0.0, 1.3)
Other Injury/Poisoning	14	12.0 (4.5, 16.0)	6	2.6 (0.5, 4.6)	9	2.5 (0.9, 4.2)
ENROLLED >= 24 MONTHS		(n=8589)		(n=17,319)		(n=26,620)
Total deaths	45	52.4 (37.1, 67.7)	39	22.5 (15.5, 29.6)	17	6.4 (3.4, 9.4)
Self-inflicted Injury/Poisoning	16	18.6 (9.5, 27.7)	20	11.5 (6.5, 16.6)	1	0.4 (0.0, 1.1)
Other Injury/Poisoning	11	12.8 (5.2, 20.4)	4	2.3 (0.0, 4.6)	6	2.3 (0.5, 4.1)
ENROLLED >= 36 MONTHS		(n=6071)		(n=12,627)		(n=20,594)
Total deaths	28	46.1 (29.1, 63.2)	31	24.6 (15.9, 33.2)	12	5.8 (2.5, 9.1)
Self-inflicted Injury/Poisoning	10	16.5 (6.3, 26.7)	14	11.1 (5.3, 16.9)	0	0.0 (0.0, 1.5)
Other Injury/Poisoning	8	13.2 (4.1, 22.3)	4	3.2 (0.1, 6.3)	4	1.9 (0.0, 3.8)